



3415A 3 Ave NW, Calgary, Alberta, T2N 0M4, Canada

**Patient Name:** Patient, Name**Specimen ID (SID):** 25-001-0000**External SID:** 123456789**Specimen Type:** Plasma**DOB:** 01-Jan-2000**Doctor:** Dr. Doctor**Date/Time Collected:** 01-Jan-2025 / 00:00**PHN:** AB 0000000**Report Date:** 12-Mar-2025**Specimen Source:** MitogenDx**Reason for Testing:** Autoinflammatory syndrome**Relevant Medications:** -**Cytokine, Chemokine & Growth Factor Panel****Laboratory Developed Test (LDT)****Report Summary:****Sample Comments:**

-

Results Summary:

High Analytes: EGF, FGF-2, FLT-3L, Fractalkine, G-CSF, GM-CSF, IFN- α 2, IFN γ , IL-1 α , IL-1 β , IL-1RA, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12p40, IL-12p70, IL-13, IL-15, IL-17A, IL-17E/IL-25, IL-17F, IL-22, IL-29, IL-33, MCP-3, M-CSF, MDC, MIP-1 α , sCD40L, TGF α , TNF α , TNF β , VEGF-A

Moderate High Analytes: GRO α , sCD137

Moderate Low Analytes: SDF-1

Low Analytes: MIP-1 δ , SCF

Results Interpretation:

The grouping profile indicates significant elevations in group A1, which may reflect innate or autoimmune inflammation, group A2, suggesting strong pro-inflammatory cytokine activity and active T helper cell-mediated inflammation, and group A3, which could suggest an expansion of immune cells.

The immune activation profile reveals an active type 1 response characterized by initiation/propagation factors (IL-12p70, IL-2) and effector cytokines (IFN γ , TNF α , TNF β), an active type 2 response with initiation/propagation factors (IL-4, IL-17E/IL-25, IL-33) and effector cytokines (IL-5, IL-13, IL-9), and a potential pathogenic type 3 response indicated by effector cytokines (IL-17A, IL-17F, IL-22) and pathogenic Th17 markers (IFN γ , GM-CSF). Additionally, there is an active anti-inflammatory profile with high levels of IL-10 and IL-1RA.

The cytokine profile suggests potential involvement of various immune cells, including B cells (IL-4, IL-6, IL-7, sCD40L), T cells (IL-15, IL-2, IL-7), cytotoxic T cells (GM-CSF, IL-12p70, IL-2, TNF α), NK cells (IL-12p70, IL-15, IL-2, TNF β), neutrophils (G-CSF, GM-CSF, IL-6, IL-8), basophils (GM-CSF, IL-3, IL-33, IL-4), dendritic cells (FLT-3L, GM-CSF, IL-3), mast cells (IL-33, IL-4, IL-9), and eosinophils (GM-CSF, IL-3, IL-5).

Disclaimer:

The interpretation of these test results should be correlated with clinical findings and other diagnostic tests. Biomarker levels can vary due to many biological, physiological, and diurnal factors; their clinical significance must be assessed by a qualified healthcare professional. This information is not intended to be used as the sole basis for diagnosis or treatment decisions.

Reviewed by: DP**Eve Technologies Corporation is a CLIA certified High Complexity International Laboratory**



3415A 3 Ave NW, Calgary, Alberta, T2N 0M4, Canada

**Patient, Name**

PHN: AB 0000000; DOB: 01-Jan-2000; SID: 25-001-0000; External SID: 123456789

Cytokine, Chemokine & Growth Factor Panel**Laboratory Developed Test (LDT)**

Analyte	Results (pg/ml)	Reference Interval†	Analyte	Results (pg/ml)	Reference Interval†
6CKine	812	293 - 1243	IL-20	32.4	5.7 - 99.9
APRIL	290	52 - 1476	IL-21	8.9	0 - 22.0
BAFF	459	285 - 1689	IL-22	492 HIGH	0 - 148
BCA-1	52.1	15 - 168	IL-23	1547	0 - 3213
CCL28	136	0 - 574	IL-24	325	24 - 1240
CTACK	390	300 - 1415	IL-27	1789	324 - 4151
EGF	213 HIGH	0 - 78.6	IL-28A	18.6	0 - 42.5
ENA-78	224	52 - 1084	IL-29	32.9 HIGH	0 - 31.8
Eotaxin	28.1	5.5 - 48.8	IL-31	26.9	0 - 37.5
Eotaxin-2	75.4	42 - 361	IL-33	48.5 HIGH	0 - 42.0
Eotaxin-3	31.7	8.2 - 76.7	IL-34	47.0	4.9 - 82.4
FGF-2	381 HIGH	16 - 225	IL-35	198	0 - 362
FLT-3L	55.0 HIGH	0.7 - 29.0	IP-10	110	21 - 281
Fractalkine	616 HIGH	32 - 305	I-TAC	19.9	9 - 289
GCP-2	25.8	5 - 190	LIF	7.3	0 - 17.3
G-CSF	198 HIGH	0 - 81.1	Lymphotactin	38.7	0 - 85.7
GM-CSF	641 HIGH	0 - 62.6	MCP-1	246	36 - 337
Granzyme A	44.5	6 - 109	MCP-2	8.6	5.9 - 35.3
Granzyme B	9.3	0 - 40.3	MCP-3	151 HIGH	3.9 - 38.6
GRO α	35.6	0.8 - 36.0	MCP-4	19.8	16 - 148
HMGB1	908	0 - 3924	M-CSF	359 HIGH	4 - 284
I-309	6.0	0 - 33.2	MDC	1243 HIGH	94 - 1213
IFN- α 2	422 HIGH	13 - 128	MIG	1423	381 - 5907
IFN β	56.1	0 - 99.1	MIP-1 α	324 HIGH	12.1 - 93.0
IFN γ	15.7 HIGH	0.2 - 8.3	MIP-1 β	31.3	9.7 - 65.6
IFN ω	28.4	0 - 55.7	MIP-1 δ	837 LOW	862 - 4175
IL-1 α	355 HIGH	0 - 74.8	MIP-3 α	19.5	1.7 - 31.2
IL-1 β	86.6 HIGH	0 - 46.2	MIP-3 β	99.4	29 - 239
IL-1RA	77.0 HIGH	1.0 - 35.5	MPIF-1	152	20 - 547
IL-2	32.8 HIGH	0 - 7.5	PDGF-AA	648	21 - 2962
IL-3	7.0 HIGH	0 - 3.5	PDGF-AB/BB	5308	1130 - 16525
IL-4	21.9 HIGH	0 - 3.3	Perforin	4102	1600 - 10826
IL-5	19.7 HIGH	0.5 - 16.9	RANTES	844	194 - 2150
IL-6	13.4 HIGH	0.2 - 10.8	sCD137	23.0	2.1 - 25.2
IL-7	17.2 HIGH	0 - 7.5	sCD40L	2493 HIGH	21 - 1040
IL-8	41.9 HIGH	0 - 21.2	SCF	219 LOW	247 - 1820
IL-9	57.4 HIGH	0 - 22.8	SDF-1	923	849 - 2770
IL-10	24.5 HIGH	0 - 19.5	sFas (ng/ml)	4.5	2.4 - 30.6
IL-11	5.8	0 - 28.3	sFasL	140	28 - 400
IL-12p40	1305 HIGH	7 - 220	TARC	19.0	1 - 106
IL-12p70	112 HIGH	0 - 21.5	TGF α	> 1250 HIGH	0.8 - 18.7
IL-13	633 HIGH	5 - 162	TNF α	743 HIGH	11 - 107
IL-15	54.5 HIGH	1.2 - 22.3	TNF β	98.2 HIGH	0 - 27.6
IL-16	226	25 - 1033	TPO	89.5	27 - 548
IL-17A	266 HIGH	0 - 24.5	TRAIL	26.2	7.9 - 92.7
IL-17E/IL-25	4297 HIGH	35 - 1545	TSLP	1.5	0 - 2.5
IL-17F	2682 HIGH	0 - 54.0	VEGF-A	108 HIGH	0 - 91.0
IL-18	77.7	3 - 235			

† Reference intervals estimated by data-mining ≥ 2000 PLASMA samples drawn from both healthy and pathological subjects.

Patient, Name

PHN: AB 0000000; DOB: 01-Jan-2000; SID: 25-001-0000; External SID: 123456789

Cytokine Groupings - Immune Signatures

Groupings represent co-expressing cytokines identified by non-biased clustering of >130 clinical plasma-EDTA samples

Analyte	Results (pg/ml)	Reference Interval†	Analyte	Results (pg/ml)	Reference Interval†
GROUP A1 - INNATE/AUTOIMMUNE INFLAMMATION			GROUP D1 - LYMPHOCYTE RECRUITMENT/ACTIVATION		
FGF-2	381 HIGH	16 - 225	BCA-1	52.1	15 - 168
IFN- α 2	422 HIGH	13 - 128	CCL28	136	0 - 574
IL-1 α	355 HIGH	0 - 74.8	I-309	6.0	0 - 33.2
IL-1 β	86.6 HIGH	0 - 46.2	IL-16	226	25 - 1033
IL-1RA	77.0 HIGH	1.0 - 35.5	IL-23	1547	0 - 3213
IL-2	32.8 HIGH	0 - 7.5	IL-35	198	0 - 362
IL-17A	266 HIGH	0 - 24.5	Lymphotactin	38.7	0 - 85.7
GROUP A2 - PRO-INFLAMMATORY/T CELL BIOMARKERS			sCD137	23.0	2.1 - 25.2
Fractalkine	616 HIGH	32 - 305	sFasL	140	28 - 400
IFN γ	15.7 HIGH	0.2 - 8.3	TPO	89.5	27 - 548
IL-4	21.9 HIGH	0 - 3.3	GROUP D2 - MUCOSAL INFLAMMATION/DAMAGE		
IL-5	19.7 HIGH	0.5 - 16.9	Eotaxin-3	31.7	8.2 - 76.7
IL-9	57.4 HIGH	0 - 22.8	HMGB1	908	0 - 3924
IL-12p40	1305 HIGH	7 - 220	IFN β	56.1	0 - 99.1
IL-12p70	112 HIGH	0 - 21.5	IFN ω	28.4	0 - 55.7
IL-13	633 HIGH	5 - 162	IL-11	5.8	0 - 28.3
IL-17F	2682 HIGH	0 - 54.0	IL-17E/IL-25	4297 HIGH	35 - 1545
IL-22	492 HIGH	0 - 148	IL-20	32.4	5.7 - 99.9
MCP-3	151 HIGH	3.9 - 38.6	IL-21	8.9	0 - 22.0
MIP-1 α	324 HIGH	12.1 - 93.0	IL-24	325	24 - 1240
TGF α	> 1250 HIGH	0.8 - 18.7	IL-28A	18.6	0 - 42.5
TNF α	743 HIGH	11 - 107	IL-29	32.9 HIGH	0 - 31.8
TNF β	98.2 HIGH	0 - 27.6	IL-31	26.9	0 - 37.5
GROUP A3 - HEMATOPOIETIC GROWTH FACTORS			IL-33	48.5 HIGH	0 - 42.0
GM-CSF	641 HIGH	0 - 62.6	IL-34	47.0	4.9 - 82.4
G-CSF	198 HIGH	0 - 81.1	LIF	7.3	0 - 17.3
IL-3	7.0 HIGH	0 - 3.5	TSLP	1.5	0 - 2.5
IL-7	17.2 HIGH	0 - 7.5	GROUP E - IMMUNE CELL TRAFFICKING/ACTIVATION		
GROUP B - INNATE INFLAMMATION/CYTOKINE STORM			6CKine	812	293 - 1243
BAFF	459	285 - 1689	CTACK	390	300 - 1415
FLT-3L	55.0 HIGH	0.7 - 29.0	Eotaxin	28.1	5.5 - 48.8
IL-27	1789	324 - 4151	Eotaxin-2	75.4	42 - 361
IL-6	13.4 HIGH	0.2 - 10.8	MDC	1243 HIGH	94 - 1213
IL-8	41.9 HIGH	0 - 21.2	MIP-1 δ	837 LOW	862 - 4175
IP-10	110	21 - 281	MPIF-1	152	20 - 547
I-TAC	19.9	9 - 289	RANTES	844	194 - 2150
IL-10	24.5 HIGH	0 - 19.5	SCF	219 LOW	247 - 1820
IL-15	54.5 HIGH	1.2 - 22.3	SDF-1	923	849 - 2770
IL-18	77.7	3 - 235	GROUP F - PLATELET ACTIVATION/WOUND HEALING		
MCP-1	246	36 - 337	APRIL	290	52 - 1476
MCP-2	8.6	5.9 - 35.3	EGF	213 HIGH	0 - 78.6
M-CSF	359 HIGH	4 - 284	ENA-78	224	52 - 1084
MIG	1423	381 - 5907	GCP-2	25.8	5 - 190
MIP-1 β	31.3	9.7 - 65.6	GRO α	35.6	0.8 - 36.0
MIP-3 α	19.5	1.7 - 31.2	MCP-4	19.8	16 - 148
MIP-3 β	99.4	29 - 239	PDGF-AA	648	21 - 2962
GROUP C - CELL DEATH BIOMARKERS			PDGF-AB/BB	5308	1130 - 16525
Perforin	4102	1600 - 10826	sCD40L	2493 HIGH	21 - 1040
sFas (ng/ml)	4.5	2.4 - 30.6	TARC	19.0	1 - 106
TRAIL	26.2	7.9 - 92.7	VEGF-A	108 HIGH	0 - 91.0

† Reference intervals estimated by data-mining ≥ 2000 PLASMA samples drawn from both healthy and pathological subjects.

Cytokine Groupings Descriptions

GROUP A1 - INNATE / AUTOIMMUNE INFLAMMATION

The analytes in this group are associated with innate immunity (IL-1 α / β , IL-17E/IL-25, IFN α 2), type 1 (IFN α 2, IL-2, MIP-1 α), and type 3 (IL-17A, IL-1) immune responses. IL-1, type I interferons, IL-17, MIP-1 α , and FGF-2 contribute to autoimmune diseases, while IL-2 and IL-17E/IL-25 can either promote or suppress autoimmunity. IL-17A and FGF-2 synergistically drive inflammation in autoimmune arthritis. IL-1, IL-17, and FGF-2 potentiate Th17-mediated immunity, a key driver of autoimmunity, whereas IL-2 and IL-17E/IL-25 negatively regulate Th17 activity. IFN α 2 exacerbates Th17-mediated inflammation, as seen in systemic lupus erythematosus (SLE), where IFN α 2 and IL-17A form a pathogenic signaling axis. IL-1 α / β also drive innate inflammatory responses and autoinflammatory conditions, and IL-1RA is expressed as a negative regulator of IL-1 signaling.

GROUP A2 - PRO-INFLAMMATORY/T CELL BIOMARKERS

This group of analytes includes pro-inflammatory cytokines involved in initiating innate inflammation and adaptive immune responses. The cytokine profile reflects Th1 (IFN γ , IL-12p70, TNF β ; intracellular pathogens/autoimmunity), Th2 (IL-4, IL-5, IL-13, IL-9; helminths/allergy/tissue repair), Th17 (IL-17F, IL-22; extracellular pathogens/autoimmunity), Th9 (IL-9), and Th22 (IL-22, IL-13) responses, which influence allergy and autoimmunity. Mixed T cell cytokine patterns may indicate diverse inflammatory responses, T cell heterogeneity and plasticity, or hybrid cells expressing multiple cytokines (e.g., IL-4 with IFN γ , IFN γ with IL-17A). These patterns may also reflect regulatory mechanisms, such as type 2 cytokine release following tissue damage from type 1 or type 3 responses.

GROUP A3 - HEMATOPOIETIC GROWTH FACTORS

The analytes in this group are hematopoietic growth factors and could indicate the expansion and activation of lymphocytes (IL-7) and/or leukocytes (GM-CSF, G-CSF, IL-3).

GROUP B - INNATE INFLAMMATION/CYTOKINE STORM

High levels of these analytes may indicate innate immune responses. IL-6 drives acute phase protein release, IL-18 acts as a pro-inflammatory alarmin via inflammasome activation, and Flt-3L supports innate lymphoid cell development. Elevated levels can signify severe systemic inflammation, such as cytokine storm (CRS). Key cytokines involved in CRS include IL-6, IL-10, IL-18, IL-8, MIG, IP-10, MIP-1 β , and MCP-1. IL-10, despite its anti-inflammatory role, is upregulated in CRS, reflecting an insufficient regulatory response. High analyte levels are common in CRS-related conditions like macrophage activation syndrome (MAS), adult-onset Still's disease (AOSD), systemic arthritis, hemophagocytic lymphohistiocytosis (HLH), and lymphocytic leukemia.

GROUP C - CELL DEATH BIOMARKERS

The analytes in this group promote cell death through facilitating (perforin) or directly inducing apoptosis (sFas, TRAIL).

GROUP D1 - LYMPHOCYTE RECRUITMENT/ACTIVATION

Elevated levels of these analytes may reflect the recruitment and activation of NK, T, and B cells. Granzyme A and B are cytotoxic mediators from NK and CD8+ T cells. sCD137 indicates NK and T cell activity, while sFasL regulates apoptosis and is shed by NK and CD8+ T cells. Lymphotactin (CD8+ T cells), CCL28 (NK and T cells), I-309 and IL-16 (CD4+ T cells) recruit lymphocytes to inflammation sites. IL-23 has context-specific pro-inflammatory effects on NK cells, CD4+ and CD8+ T cells, while IL-35 suppresses inflammation and cytotoxic cell function.

GROUP D2 - MUCOSAL INFLAMMATION/DAMAGE

Elevated levels of these analytes may indicate tissue injury and mucosal inflammation. IL-17E/IL-25, TSLP, and IL-33 are epithelial alarmins that activate type 2 immune responses. IFN β and IFN ω (type 1 interferons) and IL-28A and IL-29 (type 3 interferons) are linked to innate antiviral responses and mucosal immunity. HMGB1, released by damaged cells, promotes interferon expression. IL-34 supports mucosal-resident macrophages, while IL-11, IL-20, and IL-21 contribute to epithelial defense and tissue repair.

GROUP E - IMMUNE CELL TRAFFICKING/ACTIVATION

The analytes in this group drive the recruitment, homing and activation of leukocytes and lymphocytes.

GROUP F - PLATELET ACTIVATION/WOUND HEALING

High levels of these analytes suggest platelet activation and wound healing, as they are released by platelets and involved in angiogenesis, tissue remodeling, and inflammation. Elevated levels are seen in conditions linked to vascular injury, angiogenesis, and thrombocytosis, such as AOSD, Kawasaki disease, juvenile arthritis, FMF, COVID-19, and Crohn's disease. Lower levels are associated with thrombocytopenia-related conditions like HLH, lymphocytic leukemia, and hematopoietic stem cell transplantation. Notably, serum samples show significantly higher analyte levels than plasma samples from the same individuals.

Descriptions of the analytes and groupings with citations are available from Eve Diagnostics.

Clusters of co-expressing cytokines were determined with unsupervised clustering analysis of >130 plasma-EDTA specimens, using a similar approach as described in our publication: [Polley DJ, et al. \(2023\) Identification of novel clusters of co-expressing cytokines in a diagnostic cytokine multiplex test. Front. Immunol. 14:1223817. doi: 10.3389/fimmu.2023.1223817](#). The designations of physiological/pathological significance assigned to each grouping are speculative, based on an analysis of the immune signatures in our database of clinical specimens and on the functional/pathological roles of the analytes in each grouping established in the scientific literature.